

Durham Research Online

Deposited in DRO:

13 May 2014

Version of attached file:

Published Version

Peer-review status of attached file:

Peer-reviewed

Citation for published item:

Lyratzopoulos, G. and Abel, G.A. and McPhail, S. and Neal, R.D. and Rubin, G. (2013) 'Gender inequalities in the promptness of diagnosis of bladder and renal cancer after symptomatic presentation : evidence from secondary analysis of an English primary care audit survey.', *BMJ open.*, 3 (6). 002861.

Further information on publisher's website:

<http://dx.doi.org/10.1136/bmjopen-2013-002861>

Publisher's copyright statement:

This is an open-access article distributed under the terms of the Creative Commons Attribution Non-commercial License, which permits use, distribution, and reproduction in any medium, provided the original work is properly cited, the use is non commercial and is otherwise in compliance with the license.

Additional information:

Use policy

The full-text may be used and/or reproduced, and given to third parties in any format or medium, without prior permission or charge, for personal research or study, educational, or not-for-profit purposes provided that:

- a full bibliographic reference is made to the original source
- a [link](#) is made to the metadata record in DRO
- the full-text is not changed in any way

The full-text must not be sold in any format or medium without the formal permission of the copyright holders.

Please consult the [full DRO policy](#) for further details.

Gender inequalities in the promptness of diagnosis of bladder and renal cancer after symptomatic presentation: evidence from secondary analysis of an English primary care audit survey

Georgios Lyratzopoulos,¹ Gary A Abel,¹ Sean McPhail,² Richard D Neal,³ Gregory P Rubin⁴

To cite: Lyratzopoulos G, Abel GA, McPhail S, *et al*. Gender inequalities in the promptness of diagnosis of bladder and renal cancer after symptomatic presentation: evidence from secondary analysis of an English primary care audit survey. *BMJ Open* 2013;**3**:e002861. doi:10.1136/bmjopen-2013-002861

► Prepublication history and additional material for this paper is available online. To view these files please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2013-002861>).

Received 11 March 2013
Accepted 29 April 2013

This final article is available for use under the terms of the Creative Commons Attribution Non-Commercial 2.0 Licence; see <http://bmjopen.bmj.com>

For numbered affiliations see end of article.

Correspondence to

Dr Georgios Lyratzopoulos;
gl290@medschl.cam.ac.uk

ABSTRACT

Objectives: To explore whether women experience greater delays in the diagnosis of bladder and renal cancer when first presenting to a general practitioner with symptoms caused by those cancers and potential reasons for such gender inequalities.

Design: Prospective national audit survey of cancer diagnosis.

Setting: English primary care (2009–2010).

Participants: 920 patients with bladder and 398 patients with renal cancer (252 (27%) and 165 (42%), respectively, were women).

Primary and secondary outcome measures:

Proportion of patients with three or more pre-referral consultations; number of days from first presentation to referral; proportion of patients who presented with haematuria and proportion of patients investigated in primary care.

Results: Women required three or more prereferral consultations more often than men (27% (95% CI 21% to 33%) vs 11% (9% to 14%) for bladder ($p<0.001$); and 30% (22% to 39%) vs 18% (13% to 25%) for renal cancer ($p=0.025$)) and had a greater number of days from presentation to referral. In multivariable analysis (adjusting for age, haematuria status and use of primary care-led investigations), being a woman was independently associated with higher odds of three or more pre-referral consultations (OR=3.29 (2.06 to 5.25, $p<0.001$) for bladder cancer; and OR=1.90 (1.06 to 3.42, $p=0.031$) for renal cancer). Although presentation with haematuria was associated with more timely diagnosis of bladder cancer, gender inequalities did not vary by haematuria status for either cancer ($p=0.18$ for bladder and $p=0.27$ for renal). Each year in the UK, approximately 700 women with either bladder or renal cancer experience a delayed diagnosis because of their gender, of whom more than a quarter (197, or 28%) present with haematuria.

Conclusions: There are notable gender inequalities in the timeliness of diagnosis of urological cancers. There is a need to both reinforce existing guidelines on haematuria investigation and develop new

ARTICLE SUMMARY

Article focus

- Limited previous evidence suggests that women with urinary tract cancers may be diagnosed less promptly than men with the same cancers.
- Evidence is needed from contemporary clinical data sources to establish whether gender inequalities do exist, their magnitude, and their potential causes.

Key messages

- Women with bladder and renal cancer are more likely than men to require three or more prereferral consultations with a general practitioner, and to experience longer time intervals between presentation and hospital referral.
- There were gender differences for patients both with and without haematuria, suggesting that doctors often interpret the clinical importance of haematuria differently in men and women.
- Population health impact estimates suggest that gender inequalities can be reduced by reinforcing existing clinical guidelines on haematuria management. However, new approaches (such as use of clinical decision support tools) also need to be developed to improve the diagnosis of patients of either gender who present without haematuria.

Strengths and limitations of this study

- We were able to explore potential confounding of gender differences in diagnosis by gender differences in the management of haematuria or by differential use of primary care-led investigations.
- We have estimated the population health impact of the observed relative differences in the timeliness of diagnosis.
- The sample size of the study was relatively small (particularly for patients with renal cancer).

diagnostic decision aids and tests for patients who present without haematuria.

Gender inequalities in diagnosis of urinary tract cancer

INTRODUCTION

Promptly diagnosing patients who present with symptoms caused by cancer is a pressing priority for health-care systems worldwide.^{1–5} Globally, most patients with cancer first present to a non-specialist doctor (usually a general practitioner), who has to appropriately suspect the diagnosis in order to instigate an onward referral to a specialist. Some diagnostic delays therefore occur after presentation to a general practitioner,⁶ because signs and symptoms are initially attributed to a benign cause. Patients with certain sociodemographic characteristics may be at higher risk of experiencing a less prompt specialist referral.^{7–8} Avoiding delays in diagnosis after presentation to a doctor is an important determinant of patient experience and matters greatly to all patients and their carers.^{9–12}

In England, uniquely pronounced gender inequalities in relative survival from bladder cancer exist, with 5-year relative survival for men being 57%, compared with 44% for women.¹³ Further, specifically for bladder and renal cancer (and using English patient-reported data), notable gender inequalities have been reported in the number of times patients had to see their general practitioner with cancer symptoms before referral to a specialist (table 1).⁸ A US study also showed that women presenting with haematuria to a primary care physician are referred to a urologist for investigation less promptly than men.¹⁴ Misattribution of symptoms of urinary tract cancers in women to benign urogenital causes (eg, urinary tract infection) has been hypothesised as the reason for these gender differences,⁸ but several uncertainties remain: are gender differences in the number of pre-referral consultations apparent when using data sources other than patient surveys? Do differences in the number of prereferral consultations translate to differences in the time interval between first presentation and specialist referral? Could at least some of the gender differences in the promptness of diagnosis be explained by differences in presenting signs and symptoms (particularly haematuria)? Against this background, we set out to examine gender differences in the promptness of diagnosis of bladder and renal cancer.

METHODS

Data

We analysed data from the (English) National Audit of Cancer Diagnosis in Primary Care (2009–2010).¹⁵ Data on different aspects of the diagnostic process were collected by general practitioners or other primary care professionals in an estimated total of 1170 general practices (~14% of all practices in England). Details of the methods used in the audit have been published previously.^{15–16} Although general practice participation was voluntary, comparisons with cancer registration statistics indicate that the data set is representative of the age, sex and cancer type breakdown of patients with cancer in England.¹⁵ Data were available on two interrelated measures of promptness of diagnosis,¹⁶ that is, the number of prereferral consultations and the primary care interval (ie, the time interval between the first symptomatic presentation of a patient with cancer to a general practitioner and their first specialist referral for further investigation,⁶ measured in days by subtracting the date of first hospital referral from the date of first presentation). We used a binary form of the number of prereferral consultations (three or more vs one or two consultations) reflecting the use of this outcome by both patient groups and UK policy-makers for purposes of public reporting of the performance of NHS hospitals.¹⁷ Data were also available for patients' 5 year age group, sex, main presenting symptom and primary care investigations.

Analysis

Initial analysis was restricted to patients with at least one recorded general practitioner consultation before hospital referral and complete information on sex and age group.

In univariable analysis, we examined crude gender differences in promptness of diagnosis. We also described gender differences in haematuria status and in use of 'blood test' and ultrasound scan investigations (as these three factors have the potential to at least partially explain gender differences in promptness of diagnosis). Symptom analysis was restricted to macroscopic haematuria because it was the most common main presenting

Table 1 Gender differences in the promptness of diagnosis of urinary tract cancer (Cancer Patient Experience Survey, 2010)⁸

	Patients who saw their GP with cancer symptoms (N)	Patients with three or more GP consultations before hospital referral (n)	Patients with three or more GP consultations before hospital referral (n/N=%)	Crude odds ratio (95% confidence interval)	Adjusted odds ratio (95% confidence interval)
Bladder					
Men	4254	624	14.7	Reference	Reference
Women	1295	357	27.6	2.29 (1.97 to 2.67)	2.33 (1.99 to 2.72)
Renal					
Men	391	103	26.3	Reference	Reference
Women	208	77	37.0	1.63 (1.12 to 2.36)	1.65 (1.13 to 2.40)

GP, general practitioner.

symptom, because of its singularly strong association with urological cancers^{18–22} and because of the UK clinical guidelines by the National Institute of Health and Clinical Excellence (NICE) mandating urgent specialist referral of patients who present with macroscopic haematuria, independently of their gender ('male or female adult patients of any age who present with painless macroscopic haematuria should be referred urgently').²³ Hereafter, in this manuscript and tables, the term 'haematuria' will denote macroscopic haematuria. Investigations were restricted to 'blood test' (not otherwise specified in the data set) and ultrasound scan because they were the only two commonly recorded investigations (see online supplementary appendix 1).

Using multivariable logistic regression, we further explored the association between the number of prereferral consultations (three or more consultations vs one or two) and gender. First, we examined to what extent gender differences are confounded by other variables (age, haematuria status, ultrasound scan investigation, and 'blood test'). Subsequently, we used a final ('full') model including all variables. We further explored interactions of gender by all other variables. Because patients attend different practices, we used a sandwich estimator of standard errors.

Sensitivity analysis

We first repeated the multivariable regression model using different definitions of binary categories of the number of prereferral consultations (two or more vs one; and four or more vs one, two or three consultations). We further explored potential bias arising from missing data, using multiple imputation to produce a complete data set,^{24–27} and repeated the multivariable analysis (outlined above). Multiple imputation assumes that data are 'missing at random' (MAR), that is, that any systematic differences between the missing and observed values can be estimated using information from the observed data. The imputation model included all variables used in the analysis model and, in addition, the primary care interval variable (as the primary care interval is strongly correlated with the proportion of three or more prereferral consultations¹⁶).

Population health impact

Assuming that the national audit data are generalisable to contemporary UK practice, we illustrate the potential

population health impact of the findings. For this estimation, we used a logistic regression model (three or more vs one or two consultations) including age, gender and haematuria status as covariates to predict the proportion of men and women (with and without haematuria) who require three or more prereferral consultations both in the presence and absence of gender differences. When assuming no gender differences, we replace the odds ratio for gender by 1. The obtained proportions are scaled using national incidence statistics to give annual numbers for the UK. Stata V.11 was used for all analyses, including the uses of *ice* and *mim* commands for multiple imputation.²⁸

RESULTS

In total, there were 920 patients with bladder cancer and 398 patients with renal cancer, of whom 252 (27%) and 165 (42%), respectively, were women—proportions similar to contemporary UK population-based incidence statistics (28% and 38%, respectively, for bladder and renal cancer).²⁹ Unless where noted, after exclusion of patients whose diagnosis did not involve a prereferral consultation and missing data (see online supplementary appendix 2), complete case analysis was restricted to 740 patients with bladder cancer and 287 patients with renal cancer with at least one known prereferral consultation. Among patients with bladder and renal cancer, respectively, 91% and 85% were 55 years of age or older (ie, older than the age by which most women would have experienced their menopause) with no evidence of gender differences in these proportions ($p=0.77$ and $p=0.76$, respectively, for either cancer).

Univariable analysis

Women had longer primary care intervals compared with men for both cancers (table 2) and required three or more prereferral consultations more often than men for either bladder (27% vs 11%, $p<0.001$) or renal cancer (30% vs 18%, $p=0.025$, table 3). Although gender differences in the median primary care interval were relatively small (6 vs 4 days for bladder cancer and 16 vs 10 days for renal cancer), substantial differences existed in the tails of the distribution. For either bladder or renal cancer, the 75th centile in women was longer than that in men by about 2 weeks (table 2) and at the 90th centile, the difference increased to over 2 months

Table 2 Centiles of the primary care interval by gender for bladder and renal patients with cancer

Centile	Bladder cancer (n=721)			Renal cancer (n=271)		
	Men (n=525) (days)	Women (n=196) (days)	p Value	Men (n=160) (days)	Women (n=111) (days)	p Value
25th	0	0	0.0059	0	3	0.016
50th	4	6		10	16	
75th	15	32.5		31	46	
90th	39	103		64	82	

Gender inequalities in diagnosis of urinary tract cancer

Table 3 Univariable associations between gender and number of prereferral consultations, recorded haematuria, investigation by 'blood test' and ultrasound scan

	Bladder cancer (n=740)					Renal cancer (n=287)				
	Men (n=538)		Women (n=202)		p Value	Men (n=169)		Women (n=118)		p Value
	n	Per cent	n	Per cent		n	Per cent	n	Per cent	
Number of prereferral consultations										
1	320	59.5	102	50.5	<0.001	89	52.7	45	38.1	0.093
2	158	29.4	46	22.8		49	29.0	38	32.2	
3	40	7.4	23	11.4		17	10.1	16	13.6	
4	8	1.5	11	5.4		5	3.0	8	6.8	
5+	12	2.2	20	9.9		9	5.3	11	9.3	
1–2	478	88.8	148	73.3	<0.001	138	81.7	83	70.3	0.025
3+	60	11.2	54	26.7		31	18.3	35	29.7	
Haematuria										
Yes	394	73.2	143	70.8	0.51	57	33.7	23	19.5	0.008
No	144	26.8	59	29.2		112	66.3	95	80.5	
Ultrasound scan										
Yes	39	7.2	36	17.8	<0.001	44	26.0	35	29.7	0.50
No	499	92.8	166	82.2		125	74.0	83	70.3	
Blood test										
Yes	207	38.5	47	23.3	<0.001	71	42.0	45	38.1	0.510
No	331	61.5	155	76.7		98	58.0	73	61.0	

for bladder cancer and to about 3 weeks for renal cancer.

About two-thirds of all patients with bladder cancer and about one-quarter of all patients with renal cancer had haematuria as the main recorded primary symptom (table 3)—proportions similar to those previously reported in relevant patient populations (see online supplementary appendix 3).^{18 19 21}

Multivariable analysis

Haematuria status was strongly related to having three or more prereferral consultations but explained only a small amount of the crude gender difference for either cancer (table 4 and online supplementary appendix 4). Similarly, there was no or a limited degree of confounding of the gender association by the other three variables (see online supplementary appendix 4). Using the full model, for bladder cancer, we find that three or more prereferral consultations were substantially more likely in women (OR=3.29 (2.06 to 5.25, $p<0.001$, table 4)) and less likely among those presenting with haematuria (OR=0.29 (0.19 to 0.46, $p<0.001$). For renal cancer, three or more prereferral consultations were more likely in women (OR=1.90 (1.06 to 3.42, $p=0.031$)) without evidence for an association with haematuria ($p=0.25$).

Regarding interactions, there was no evidence that gender differences in the proportion of patients with three or more prereferral consultations varied between patients with and without haematuria ($p=0.18$ for bladder and $p=0.27$ for renal cancer), as well as by age group ($p=0.38$ and $p=0.10$), 'blood test' use ($p=0.71$ and $p=0.91$) or ultrasound scan use ($p=0.20$ and $p=0.59$).

Sensitivity analysis

Repeating the multiple logistic regression using different binary categories of the number of prereferral consultations produced similar findings, particularly regarding associations with gender and haematuria status. Sensitivity analysis using multiple imputation of missing data also produced similar findings (see online supplementary appendices 5 and 6).

Population health impact

Each year in the UK, about 2900 women are diagnosed with bladder cancer and 3000 with renal cancer.²⁹ Of those women, we estimate that each year approximately 693 (~435 with bladder cancer and ~258 with renal cancer) experience three or more prereferral consultations when they would have required only one or two had gender inequalities not been present (see online supplementary appendix 7). More than a quarter (197, or 28%) of those women experience gender inequalities in the presence of haematuria.

DISCUSSION

In our study population, we found that women were more likely than men to experience a non-prompt diagnosis of bladder and renal cancer. Being a woman was an independent risk factor for a less timely diagnosis even after adjustment for age and presence/absence of haematuria or use of investigations. Moreover, differences in haematuria status between men and women explain only a small fraction of the crude gender differences. The findings indicate that generalists are less likely to suspect the diagnosis of urinary tract cancers in

Gender inequalities in diagnosis of urinary tract cancer

Table 4 Crude associations and independent predictors of three or more prereferral consultations from the 'full' model (adjusted for gender, age, haematuria status and investigation status)

	Crude odds ratio	95% UCL	95% LCL	p Value*	Adjusted odds ratio	95% UCL	95% LCL	p Value*
Bladder (n=740)								
Men	Reference			<0.001	Reference			<0.001
Women	2.91	1.93	4.39		3.29	2.06	5.25	
16–54	1.52	0.75	3.10	0.23	1.20	0.53	2.72	0.34
55–64	0.68	0.34	1.37		0.59	0.29	1.21	
65–74	Reference				Reference			
75–84	1.32	0.79	2.19		1.18	0.69	2.03	
85+	1.42	0.75	2.66		1.27	0.65	2.49	
No haematuria	Reference			<0.001	Reference			<0.001
Haematuria	0.28	0.18	0.42		0.29	0.19	0.46	
No blood test	Reference			<0.001	Reference			<0.001
Blood test	2.09	1.39	3.13		2.47	1.58	3.86	
No ultrasound scan	Reference			<0.001	Reference			0.18
Ultrasound scan	2.59	1.50	4.45		1.55	0.82	2.93	
Renal (n=287)								
Men	Reference			0.026	Reference			0.031
Women	1.88	1.08	3.27		1.90	1.06	3.42	
16–54	1.17	0.49	2.76	0.99	1.05	0.41	2.74	0.99
55–64	1.04	0.49	2.22		0.85	0.39	1.85	
65–74	Reference				Reference			
75–84	0.95	0.42	2.13		0.95	0.38	2.38	
85+	1.23	0.45	3.36		0.97	0.34	2.77	
No haematuria	Reference			0.023	Reference			0.25
Haematuria	0.44	0.22	0.89		0.64	0.30	1.37	
No blood test	Reference			0.001	Reference			<0.001
Blood test	2.70	1.54	4.75		2.99	1.64	5.46	
No ultrasound scan	Reference			0.015	Reference			0.023
Ultrasound scan	2.06	1.15	3.69		2.17	1.11	4.24	

*From Wald tests, with joint tests used where applicable.
LCL, lower confidence limit; UCL, upper confidence limit.

women and that haematuria is often interpreted differently by general practitioners depending on the patient's gender. Optimising referral decisions for women with bladder and renal cancer who present with haematuria may have a notable impact on gender inequalities in promptness of diagnosis. However, many patients with urinary tract cancers (both women and men) present without haematuria. For those patients, new approaches are needed to help improve the promptness of diagnosis for patients of either gender.

Our findings amplify previous limited evidence on gender inequalities in the diagnosis of urinary tract cancers using patient-reported data in England and also previous US research.^{8–14} The strengths of the present study include the use of two different measures of promptness of diagnosis; the adjustment of the analysis for haematuria status and investigation use; the examination of potential interactions between gender and all other variables and the use of sensitivity analysis (including for missing data). Considering generalisability, the sample of patients was similar to population-based incidence data in respect of age, sex and cancer,¹⁵ and to other primary care study populations in respect of the proportions of patients who presented with haematuria (see online

supplementary appendix 3).^{18–21} Further, in supplementary analysis, we compared the characteristics of 535 participating with 2349 non-participating practices and found trivial differences in practice care quality and patient experience measures, and small differences for practice population and team size (see online supplementary appendix 8).

We believe that the principal reason for improving the timeliness of cancer diagnosis among symptomatic patients is to ensure as positive an experience of cancer care as possible for all patients,^{9–12} although achieving such improvements may also help to improve treatment and prognosis for some. Indeed, there is some evidence indicating an association between delay and worse oncological outcomes for patients with bladder cancer presenting with haematuria.^{30–31} For bladder cancer in particular, women are more likely than men to be diagnosed at an advanced stage,³² and there are also uniquely large gender inequalities in relative survival, as women have a substantially worse 5-year relative survival than men (57% vs 44%, respectively).¹³ While the median primary care intervals for men and women with bladder cancer were similar (4 and 6 days, respectively), there were substantial differences in the tails of the

Gender inequalities in diagnosis of urinary tract cancer

Table 5 Comparison of survey design and methodologies used by the National Audit of Cancer Diagnosis in Primary Care,¹⁵ and the Cancer Patient Experience Survey 2010⁸

	National audit of cancer diagnosis in primary care	Cancer patient experience survey
Population	Patients with cancer in participating practices (about 14% of all English practices)	Patients with cancer across England
Sampling frame	Patients with new diagnosis of cancer during audited period (2009–2010)	Patients attending an NHS hospital for cancer treatment Jan–Mar 2010 (many of these cases have been diagnosed before the survey period)
Sample representativeness	Aimed to include all patients eligible for inclusion in the audit with minimal attrition (independently of survival length)	Only about two-thirds (67%) of all patients in the original sampling frame participated in the survey—after exclusion of people who had died, that is, patients with only short-term survival were likely to be a priori excluded
Method of outcome ascertainment	Case note review by general practitioner or other primary care professional (eg, practice nurse)	Patients responding to a specific survey question

NHS, National Health Service.

distributions that may have contributed to gender inequalities in stage and survival, although other factors such as gender differences in tumour subtypes or other biological factors may also be responsible.³³ These considerations however apply only to bladder cancer; we are not aware of evidence suggesting substantial gender differences in stage and survival from renal cancer.¹³ The findings concord with and substantially amplify previous evidence from patient survey data indicating notable inequalities in the promptness of diagnosis of women with bladder and renal cancer (table 1).⁸ The degree of concordance of the observed gender inequalities between national audit and patient-reported data is remarkable, given the differences in the methodologies used (see table 5).

The findings signal a large potential for improving the timeliness of diagnosis of urinary tract cancer in women. In part, this can be achieved by reinforcement and more rigorous adherence to existing clinical guidelines on investigation and management of haematuria.²³ Previous US research indicates that primary care physicians often do not adhere to guidelines for prompt investigation of patients presenting with haematuria.³⁴ Interventions to help general practitioners avoid initial misattribution of haematuria in women with urinary tract cancer to benign causes need to be promptly developed and evaluated. Elimination of gender differences in referral decisions in the presence of haematuria would only partially address overall gender inequalities, although it can help produce notable improvements with relative ease and speed. However, because many patients with urinary tract cancers present without haematuria, research to better understand these less specific presentations is also required. The implementation of clinical decision support tools (ideally combined with evaluation of their utility) may be helpful, since they incorporate information from multiple symptoms, signs and systemic manifestations (such as thrombocytosis).^{20–35} The application

of consultation techniques, such as ‘safety-netting’, the development of new service models such as outreach facilities for specialist consultation with urologists and the development of new tests (eg, based on biomarkers) are also worthy of exploration and prioritisation of research investment. Improvements in the sensitivity of generalist consultations to suspect the presence of urological cancer among symptomatic patients should reduce gender inequalities, but should also benefit all patients of either gender. The positive predictive value of haematuria for urological cancer is generally lower than 15% (depending on age),²⁰ and is lower in women than men.²¹ This means that even in a hypothetical situation where all patients presenting to general practitioners with haematuria as the main presenting symptom were referred promptly for specialist investigation, the great majority of them would be found not to have cancer. Nevertheless, clinical guidelines, such as those produced by NICE, mandate the referral of all patients who present with painless macroscopic haematuria independently of their gender—see also Methods.²³ Health economics analyses to explore the cost-effectiveness of these clinical protocols may be justified. These realisations can serve as potent reminders of the need for the development of newer tests (particularly easily accessible and acceptable point-of-care tests) and service models.

In conclusion, we report compelling evidence that in the study setting and during the study period women with urinary tract cancers were likely to experience a delayed diagnosis compared with men with the same cancers. Reinforcing existing guidelines on haematuria investigation and development of new diagnostic aids for patients who present without haematuria are needed. The findings should inform similar investigations in other country populations.

Author affiliations

¹Department of Public Health and Primary Care, Cambridge Centre for Health Services Research, University of Cambridge, Cambridge, UK

²National Cancer Intelligence Network (NCIN), London, UK

³North Wales Centre for Primary Care Research, College of Health and Behavioural Sciences, Bangor University, Wrexham, UK

⁴Wolfson Research Institute, School of Medicine and Health, University of Durham, Queen's Campus, University Boulevard, Stockton-on-Tees, UK

Acknowledgements We are grateful to all primary care professionals in participating practices for collecting, collating and submitting anonymous data; and respective Cancer Networks, the Royal College of General Practitioners, the National Cancer Action Team for supporting the audit and the National Clinical Intelligence Network (NCIN).

Contributors The study was conceived and the methods developed in discussions between all authors. The idea for the study is rooted in the findings of a previous collaborative project involving four of the authors using patient experience data. GL, GAA and SMP analysed the data and the findings were interpreted by all authors. In addition, RDN and GPR provided expert clinical commenting. GL wrote the first draft, which was edited over several versions by all other authors. Authors communicated frequently during the course of the project. All authors had full access to all of the data (including statistical reports and tables) in the study and can take responsibility for the integrity of the data and the accuracy of the data analysis. GL is the guarantor.

Funding The paper is independent research arising from a Post-Doctoral Fellowship award to GL supported by the National Institute for Health Research (PDF-2011-04-047). The views expressed in this publication are those of the authors and not necessarily those of the NHS, the National Institute for Health Research or the Department of Health.

Competing interests None.

Provenance and peer review Not commissioned; externally peer reviewed.

Data sharing statement Please contact the corresponding author.

REFERENCES

- Department of Health. Improving outcomes: a strategy for cancer. 2011. http://www.dh.gov.uk/en/Publicationsandstatistics/Publications/PublicationsPolicyAndGuidance/DH_123371 (accessed Apr 2013).
- Prades J, Espinas JA, Font R, *et al.* Implementing a cancer fast-track programme between primary and secondary care in Catalonia (Spain): a mixed methods study. *Br J Cancer* 2011;105:753–9.
- New South Wales Government. Cancer Plan 2011–15. 2011. <http://www.cancerplan.cancerinstitute.org.au/> (accessed Apr 2013).
- Olesen F, Hansen RP, Vedsted P. Delay in diagnosis: the experience in Denmark. *Br J Cancer* 2009;101(Suppl 2):S5–8.
- Cancer Care Ontario. Diagnostic assessment programs: an environmental scan. 2009. <https://www.cancercare.on.ca/common/pages/UserFile.aspx?fileId=64068> (accessed Apr 2013).
- Weller D, Vedsted P, Rubin G, *et al.* The Aarhus statement: improving design and reporting of studies on early cancer diagnosis. *Br J Cancer* 2012;106:1262–7.
- Neal RD, Allgar VL. Sociodemographic factors and delays in the diagnosis of six cancers: analysis of data from the 'National Survey of NHS Patients: Cancer'. *Br J Cancer* 2005;92:1971–5.
- Lyratzopoulos G, Neal RD, Barbiere JM, *et al.* Variation in number of general practitioner consultations before hospital referral for cancer: findings from the 2010 National Cancer Patient Experience Survey in England. *Lancet Oncol* 2012;13:353–65.
- Rarer Cancer Foundation. Primary cause? An audit of the experience in primary care of rarer cancer patients. 2011. <http://www.rarercancers.org.uk/images/stories/cdf/p8and9/primary%20cause%20-%20final.pdf> (accessed Apr 2013).
- Pancreatic Cancer UK. Study for survival 2011. 2011. http://www.pancreaticcancer.org.uk/media/100292/report_final_for_web.pdf (accessed Apr 2013).
- The Roy Castle Lung Cancer Foundation. Explaining variations in lung cancer in England. 2011. <http://www.roycastle.org/Resources/Roy%20Castle/Documents/PDF/Variations%20in%20lung%20cancer%20in%20England.pdf> (accessed Apr 2013).
- Teenage Cancer Trust. One in four teenage cancer patients forced to visit GP at least four times before referral. 2012. <http://www.teenagecancertrust.org/who-we-are/media-centre/press-releases/one-in-four-teenage-cancer-patients-forced-to-visit/> (accessed Apr 2013).
- Rachet B, Maringe C, Nur U, *et al.* Population-based cancer survival trends in England and Wales up to 2007: an assessment of the NHS cancer plan for England. *Lancet Oncol* 2009;10:351–69.
- Johnson EK, Daignault S, Zhang Y, *et al.* Patterns of hematuria referral to urologists: does a gender disparity exist? *Urology* 2008;72:498–502.
- Rubin G, Elliott K, McPhail S, Royal College of General Practitioners. National audit of cancer diagnosis in primary care. 2011. http://www.rcgp.org.uk/pdf/National_Audit_of_Cancer_Diagnosis_in_Primary-Care.pdf (accessed Apr 2013).
- Lyratzopoulos G, Abel GA, McPhail S, *et al.* Measures of promptness of cancer diagnosis in primary care: secondary analysis of national audit data on patients with 18 common and rarer cancers. *Br J Cancer* 2013;108:1550–1.
- Department of Health. National Cancer Patient Experience Survey 2011/12—National Report. <http://www.quality-health.co.uk/images/stories/pdfs/2012CancerReports/2011-12cancerpatientexperiencesurveynationalreport.pdf> (accessed Apr 2013).
- Shephard EA, Stapley S, Neal RD, *et al.* Clinical features of bladder cancer in primary care. *Br J Gen Pract* 2012;62:e598–604.
- Bruyninckx R, Buntinx F, Aertgeerts B, *et al.* The diagnostic value of macroscopic haematuria for the diagnosis of urological cancer in general practice. *Br J Gen Pract* 2003;53:31–5.
- Hippisley-Cox J, Coupland C. Identifying patients with suspected renal tract cancer in primary care: derivation and validation of an algorithm. *Br J Gen Pract* 2012;62:e251–60.
- Jones R, Latinovic R, Charlton J, *et al.* Alarm symptoms in early diagnosis of cancer in primary care: cohort study using General Practice Research Database. *BMJ* 2007;334:1040.
- Shephard E, Neal R, Rose P, *et al.* Clinical features of kidney cancer in primary care: a case-control study using primary care records. *Br J Gen Pract* 2013;63:250–5.
- National Institute for Health and Clinical Excellence. Referral for suspected cancer: a clinical practice guideline. London. 2005. Clinical guideline 27. <http://www.nice.org.uk/CG027> (accessed Apr 2013).
- He Y, Yucel R, Zaslavsky AM. Misreporting, missing data, and multiple imputation: improving accuracy of cancer registry databases. *Chance (NY)* 2008;21:55–8.
- Nur U, Shack LG, Rachet B, *et al.* Modelling relative survival in the presence of incomplete data: a tutorial. *Int J Epidemiol* 2010;39:118–28.
- Lyratzopoulos G, Abel GA, Barbiere JM, *et al.* Variation in advanced stage at diagnosis of lung and female breast cancer in an English region 2006–2009. *Br J Cancer* 2012;106:1068–75.
- Lyratzopoulos G, Abel GA, Brown CH, *et al.* Socio-demographic inequalities in stage of cancer diagnosis: evidence from patients with female breast, lung, colon, rectal, prostate, renal, bladder, melanoma, ovarian and endometrial cancer. *Ann Oncol* 2013;24:843–50.
- Royston P. Multiple imputation of missing values: further update of ice, with an emphasis on interval censoring. *Stata J* 2007;7:445–64.
- Office for National Statistics. Cancer incidence and mortality in the United Kingdom, 2007–2009. <http://www.ons.gov.uk/ons/publications/re-reference-tables.html?edition=tcm%3A77-251462> (accessed Apr 2013).
- Wallace DM, Bryan RT, Dunn JA, *et al.* West Midlands Urological Research Group. Delay and survival in bladder cancer. *BJU Int* 2002;89:868–78.
- Hollenbeck BK, Dunn RL, Ye Z, *et al.* Delays in diagnosis and bladder cancer mortality. *Cancer* 2010;116:5235–42.
- Barbiere JM, Saeb-Parsy K, Greenberg DC, *et al.* Trends in the use of radiotherapy and radical surgery for patients with bladder urothelial cell carcinoma in East Anglia, 1995–2006. *BJU Int* 2011;108:1106–14.
- Noon AP, Albertsen PC, Thomas F, *et al.* Competing mortality in patients diagnosed with bladder cancer: evidence of undertreatment in the elderly and female patients. *Br J Cancer* 2013;108:1534–40.
- Nieder AM, Lotan Y, Nuss GR, *et al.* Are patients with hematuria appropriately referred to Urology? A multi-institutional questionnaire based survey. *Urol Oncol* 2010;28:500–3.
- Hamilton W, Green T, Martins T, *et al.* Evaluation of risk assessment tools for suspected cancer in general practice. *Br J Gen Pract* 2013;63:30–6.

BMJ Open

Gender inequalities in the promptness of diagnosis of bladder and renal cancer after symptomatic presentation: evidence from secondary analysis of an English primary care audit survey

Georgios Lyratzopoulos, Gary A Abel, Sean McPhail, et al.

BMJ Open 2013 3:

doi: 10.1136/bmjopen-2013-002861

Updated information and services can be found at:

<http://bmjopen.bmj.com/content/3/6/e002861.full.html>

These include:

Data Supplement

"Supplementary Data"

<http://bmjopen.bmj.com/content/suppl/2013/06/04/bmjopen-2013-002861.DC1.html>

References

This article cites 24 articles, 6 of which can be accessed free at:

<http://bmjopen.bmj.com/content/3/6/e002861.full.html#ref-list-1>

Article cited in:

<http://bmjopen.bmj.com/content/3/6/e002861.full.html#related-urls>

Open Access

This is an open-access article distributed under the terms of the Creative Commons Attribution Non-commercial License, which permits use, distribution, and reproduction in any medium, provided the original work is properly cited, the use is non commercial and is otherwise in compliance with the license. See:

<http://creativecommons.org/licenses/by-nc/3.0/> and

<http://creativecommons.org/licenses/by-nc/3.0/legalcode>

Email alerting service

Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

To request permissions go to:

<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:

<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:

<http://group.bmj.com/subscribe/>

Topic Collections

Articles on similar topics can be found in the following collections

[Epidemiology](#) (684 articles)
[General practice / Family practice](#) (185 articles)
[Health services research](#) (395 articles)
[Oncology](#) (142 articles)
[Urology](#) (15 articles)

Notes

To request permissions go to:
<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:
<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:
<http://group.bmj.com/subscribe/>